



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## SUMMATION OF CERTAIN INFINITE SERIES.

By W. J. GREENSTREET, M. A., F. R. A. S., Editor of The Mathematical Gazette, Stroud, England.

I note with some surprise that no solution of Problem 221 has yet (November, 1906) reached the MONTHLY. I therefore take the liberty of pointing out the method usually adopted in questions of this type, and embody in my remarks solutions of many similar questions taken from Todhunter and Hogg's Trigonometry, Hobson's Trigonometry (our best English work on the subject), and from various Cambridge Scholarship and other papers of recent years. The whole will, I hope, form a useful summary for those to whom the methods are unfamiliar.

1. To save space, we write the series  $\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots$  to  $\infty$  in the form  $\sum_{n=1}^{n=\infty} \frac{1}{n^2}$ ;  $\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots$  to  $\infty$ , in the form  $\sum_{n=1}^{n=\infty} \frac{1}{(2n-1)^2}$ .

The sign  $\equiv$  is used for "identical with." We know that

$$\left. \begin{aligned} \frac{\sin \theta}{\theta} &= 1 - \frac{\theta^2}{3!} + \frac{\theta^4}{5!} - \frac{\theta^6}{7!} + \dots \equiv \sum_{n=1}^{n=\infty} (-1)^{n-1} \frac{\theta^{2n-2}}{(2n-1)!} \\ \text{and also } &= (1 - \frac{\theta^2}{\pi^2}) (1 - \frac{\theta^2}{2^2 \pi^2}) (1 - \frac{\theta^2}{3^2 \pi^2}) \dots \equiv \prod_{n=1}^{n=\infty} (1 - \frac{\theta^2}{n^2 \pi^2}) \end{aligned} \right\} (A).$$

Taking logarithms of the two right hand expressions, and expanding in powers of  $\theta$ , we may equate the coefficients of the respective powers of  $\theta$  in the system (A).

Thus equating coefficients of  $\theta^2$  we have

$$(a) \quad -\frac{\theta^2}{3!} = -\theta^2 \left( \sum_{n=1}^{n=\infty} \frac{1}{n^2 \pi^2} \right), \text{ whence } \sum_{n=1}^{n=\infty} \frac{1}{n^2} = \frac{\pi^2}{6}.$$

Hence, if  $a, b, c, \dots$  denote all the prime numbers except unity,

$$\begin{aligned} (b) \quad & (1 - a^{-2})^{-1} (1 - b^{-2})^{-1} (1 - c^{-2})^{-1} \dots \\ &= (1 + a^{-2} + a^{-4} + \dots) (1 + b^{-2} + b^{-4} + \dots) (1 + c^{-2} + c^{-4} + \dots) \dots \\ &= 1 + (a^{-2} + b^{-2} + c^{-2} + \dots) + \dots + (a^{-r} b^{-s} \dots)^2 + \dots \\ &= 1 + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} \dots \text{ad infinitum} = \frac{\pi^2}{6}. \end{aligned}$$

We also have at once  $\prod_2^{\infty} (1-a^{-2}) = \frac{6}{\pi^2}$ .

Again equating coefficients of  $\theta^4$  in (A), we have

$$\frac{1}{5!} - \frac{1}{2} \left( \frac{1}{3!} \right)^2 = -\frac{1}{2\pi^4} \sum_{n=1}^{\infty} \frac{1}{n^4},$$

and it follows that

$$(c) \quad \sum_{n=1}^{\infty} \frac{1}{n^4} = \frac{\pi^4}{12} \left( \frac{1}{3} - \frac{1}{5} \right) = \frac{\pi^4}{90}.$$

Then, if  $a, b, c, \dots$  are the prime numbers we have

$$\begin{aligned} (d) \quad & (1-a^{-4})^{-1}(1-b^{-4})^{-1}(1-c^{-4})^{-1}\dots \\ & = (1+a^{-4}+a^{-8}+\dots)(1+b^{-4}+b^{-8}+\dots)(1+c^{-4}+c^{-8}+\dots) \\ & = (1+a^{-4}+b^{-4}+\dots+(a^r b^s c^t \dots)^4+\dots \\ & = \sum_{n=1}^{\infty} \frac{1}{n^4} = \frac{\pi^4}{90} \text{ by (c).} \end{aligned}$$

Thus dividing (d) by (b) we have

$$\begin{aligned} (e) \quad & \prod_2^{\infty} (1+a^{-2}) = \frac{15}{\pi^2}, \text{ which is in No. 221, p. 190. And} \\ & \prod_2^{\infty} (1+a^{-2})^{-1} = \frac{2^2}{2^2+1} \cdot \frac{3^2}{3^2+1} \cdot \dots = \frac{\pi^2}{15}. \end{aligned}$$

We can connect up the series  $\sum_{n=1}^{\infty} \frac{1}{n^2}$  with  $\sum_{n=1}^{\infty} \frac{1}{(2n-1)^2}$ , as follows:

$$\text{Let } S = \frac{1}{1^n} + \frac{1}{2^n} + \frac{1}{3^n} + \dots$$

$$S_1 = \frac{1}{1^n} + \frac{1}{3^n} + \frac{1}{5^n} + \frac{1}{7^n} + \dots$$

$$\text{Then } S = \frac{1}{1^n} + \frac{1}{3^n} + \frac{1}{5^n} + \dots + \frac{1}{2^n} \left( \frac{1}{1^n} + \frac{1}{2^n} + \frac{1}{3^n} + \dots \right) = S_1 + \frac{1}{2^n}.$$

(f)  $\therefore S_1 = \frac{2^n-1}{2^n} S$ . So that if  $S$  be known,  $S_1$  is known. For instance, if  $n=2$ ,  $S = \frac{1}{6}\pi^2$  by (a),

$$(g) \quad \text{and } S_1 \equiv \sum_{n=1}^{n=\infty} \frac{1}{(2n-1)^2} = \frac{2^2-1}{2^2} S = \frac{3}{4} \cdot \frac{\pi^2}{6} = \frac{\pi^2}{8}.$$

$$\text{If } n=4, S = \frac{\pi^2}{90} \text{ by (c),}$$

$$(h) \quad \text{and } S_1 \equiv \sum_{n=1}^{n=\infty} \frac{1}{(2n-1)^4} = \frac{2^4-1}{2^4} S = \frac{15}{16} \cdot \frac{\pi^4}{90} \text{ by (c)} = \frac{\pi^4}{96}.$$

$$(i) \quad \text{Consider } \sum_{n=1}^{n=\infty} (-1)^{n-1} \sum_{n=1}^{n=\infty} \frac{1}{n^2}.$$

$$\begin{aligned} \text{Here } & \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \dots \\ &= \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots - \frac{1}{2^2} (1 + \frac{1}{2^2} + \frac{1}{3^2} + \dots) \\ &= \frac{\pi^2}{8} - \frac{\pi^2}{24} \text{ by (g) and (a)} = \frac{\pi^2}{12}. \end{aligned}$$

(j) Now consider the identity  $\left[ \sum_{n=1}^{n=\infty} \frac{1}{n^2} \right]^2 \equiv \sum_{n=1}^{n=\infty} \frac{1}{n^4} + 2 \sum \sum \frac{1}{p^2 q^2}$ , where  $p$  and  $q$  are any of the numbers 1, 2, 3, ...,  $\infty$ .

$$\text{Here } \left( \frac{\pi^2}{6} \right)^2 = \frac{\pi^4}{90} + 2 \sum \sum \frac{1}{p^2 q^2}, \text{ by (a) and (d).}$$

Then,  $\sum \frac{1}{p^2 q^2} = -\frac{\pi^4}{180} + \frac{\pi^4}{72} = \frac{\pi^4}{120}$ . Hence the sum of the series formed by multiplying together every two of the terms of the series  $\frac{1}{1^2}, \frac{1}{2^2}, \frac{1}{3^2}, \dots$ , is  $\frac{\pi^4}{120}$ .

Now consider the identity,

$$\frac{\frac{1}{2}n(n+1)}{(2n+1)^4} = \frac{1}{8} \left[ \frac{1}{(2n+1)^2} - \frac{1}{(2n+1)^4} \right].$$

$$\begin{aligned} \text{Here } \sum_{n=1}^{n=\infty} \frac{\frac{1}{2}n(n+1)}{(2n+1)^4} &= \frac{1}{8} \left[ \sum_{n=1}^{n=\infty} \frac{1}{(2n+1)^2} - \sum_{n=1}^{n=\infty} \frac{1}{(2n+1)^4} \right] \\ &= \frac{1}{8} \left[ \frac{\pi^2}{8} - \frac{\pi^4}{96} \right] \text{ by (g) and (h)} = \frac{\pi^2}{64} \left[ 1 - \frac{\pi^2}{12} \right]. \end{aligned}$$

[ To be continued. ]